



財團法人生物技術開發中心
Development Center for Biotechnology

Tri-mannosyl Antibody: A Versatile Payload ADC Platform


Development Center for Biotechnology

Contact information:
Kuan-Tsung Lin
Tel: +886-2-7700-3800#5239
E-mail: kotlin2@dcb.org.tw




Tri-mannosyl Antibody Drug Conjugation Platform





1. Site specific ADC at the N297 of an antibody.
2. Homogeneous DAR (drug antibody ratio) with **2 or 4**.
3. Homogeneous payload or dual payload: **High diversity** in cancer applications.
4. Manufacturing condition: liquid phase, room temperature, user friendly and within overnight.
5. Materials including tri-mannosyl antibody, GnT-1 and GnT-2 are **mass-produced** by cell line process developments.



Tri-mannosyl Trastuzumab-4DM1 Is Generated by Single Step Enzyme Process and SPAAC Reaction







Effective Tumor Cell Growth Inhibition by Tri-mannosyl Trastuzumab-4DM1





Coupling of First Payload DM-1 by GnT-I and SPAAC Reaction

- GlcNAc
- Mannose
- ▼ Fucose
- UDP-GlcNAz
- ◆ Payload





Coupling of The Second Payload Duocarmycin by GnT-II and SPAAC Reaction

- GlcNAc
- Mannose
- ▼ Fucose
- UDP-GlcNAz
- ◆ Payload




The Dual Payload of Tri-mannosyl Trastuzumab ADC Shows Superior Potency to Kadcyla-Resistant Breast Cancer Cell Line MDA-MB-175VII


Dual payload of tri-mannosyl ADC



Binding kinetics



MDA-MB-175VII




	Herceptin	Kadcyla	Tri-mannosyl-Trastuzumab-4GlcNAZ-2(DM1-Duocarmycin SA)
LD50 (nM)	193.42	45.72	10.03
ID50 (nM)	0.52	1.92	4.91
Net killing(%)	50.9	65.37	80.13
Kd (M)	4.23e-011	4.93e-011	3.54e-011

The Profile of Tri-mannosyl Antibody Platform


IP position	WO2018126092A and TW application number: 106146600
UDP-GlcNAz transfer percentage	Over 95%
Payload conversion rate	Over 95% for homogeneous and 90% for dual payloads
Yield	about 60% for homogeneous payload
Optimal temperature	Not determined but the room temperature is satisfied
Conversion enzyme activity	GnT-1 (limited step)= 1.5 g antibody /hr/mg (UDP-GlcNAZ as substrate)
Scale	200 mg level , 1 g level (2019)
Tri-mannosyl antibody production	CHO cell over-expressed N-acetyl glucosaminidase is in the final evaluation
GnT-1 and GnT-2 production	CHO cell over-expressed both enzymes are under way

Homogeneous Payload with DAR4



Reduced Mass

Dual Payloads with DAR4



in

37